## School Entry Requirements

Before a child enters school for the first time, they are required to receive the following shots:

DTaP (diphtheria, tetanus and acellular pertussis)	5 doses*
Polio	4 doses*
Hib (haemophilus influenzae type B)	3-4 doses**
Measles	2 doses
Mumps	2 doses
Rubella	1 dose
Hepatitis B	3 doses
Varicella (chickenpox)	2 doses <sup>+</sup>

- \* The last dose of DTaP and Polio must be given on/or after age 4.
- \*\* Children beyond their 5th birthday are not required to have any Hib vaccine.
- + Vaccination is required unless documentation of disease history is provided by a health care provider.

Note:

Pneumococcal, Hepatitis A and Flu vaccines are not required but are recommended for this age group by the Advisory Committee on Immunization Practices.

## 7<sup>th</sup> Grade Entry Vaccine Requirements:

Children entering 7<sup>th</sup> grade should be up-to-date with all of the required vaccines listed above.

In addition, all children entering 7th grade or who have reached 12 years of age on or after July 1, 2015 are required to have one dose of Tdap (tetanus, diphtheria, acellular pertussis) vaccine and one dose of meningococcal vaccine.

Note:

HPV, Hepatitis A, Flu, and a second dose of Varicella vaccine are not required but are recommended for this age group by the Advisory Committee on Immunization Practices.

## College/University Vaccine Requirements:

DTaP (diphtheria, tetanus and acellular pertussis)	3 doses <sup>1</sup>
Polio	3 doses <sup>2</sup>
Measles	2 doses <sup>3</sup>
Mumps	2 doses <sup>4</sup>
Rubella	1 dose <sup>5</sup>
Hepatitis B	3 doses <sup>6</sup>

- Footnote<sup>1</sup> Three doses of tetanus/diphtheria toxoid, of which one must be tetanus/diphtheria/pertussis if entered college or university for the first time on or after July 1, 2008.
- Footnote<sup>2</sup> An individual attending school who has attained his or her 18<sup>th</sup> birthday is not required to receive polio vaccine.
- Footnote<sup>3</sup> Measles vaccine is not required if any of the following occur: Diagnosis of disease by an MD prior to January 1, 1994; Born prior to 1957; An individual who has been documented by serological testing to have a protective antibody titer against measles. A second dose of measles vaccine is not required for individuals enrolled in college or university for the first time before July 1, 1994.
- Footnote<sup>4</sup> Mumps vaccine is not required if any of the following occur: Born prior to 1957; Enrolled in college or university before July 1, 1994; An individual who has been documented by serological testing to have a protective antibody titer against mumps. A second mumps vaccine is not required of an individual enrolled in college or university prior to July 1, 2008.
- Footnote<sup>5</sup> Rubella vaccine is not required if any of the following occur: 50 years of age or older; Enrolled in college or university before February 1, 1989 and after their 30th birthday; An individual who has been documented by serological testing to have a protective antibody titer against Rubella.
- Footnote<sup>6</sup> Hepatitis B vaccine is not required if any of the following occur: Born before July 1, 1994.
- Note: Meningococcal, HPV, Hepatitis A, Flu, and a second dose of Varicella vaccine are not required but are recommended for this age group by the Advisory Committee on Immunization Practices.



State of North Carolina
Department of Health and Human Services
www.ncdhhs.gov
N.C. DHHS is an equal opportunity employer
and provider. (Rev. 11/14)